

KALISMA



$$A = P \times \ell$$

$$= P d\theta \cdot P \sin \theta d\phi$$

$$= P^2 \sin \theta d\theta d\phi$$

$$V = P^2 \sin \theta d\theta d\phi dr$$

$$A = \int_0^{2\pi} \int_0^\pi P \sin \theta$$





$$A = r d\phi dz$$

$$V = r d\phi dz dr$$


Listrik Statis

$$q = \pm Ne$$



$$F = \frac{1}{4\pi\epsilon_0}$$

$$u_0 = \frac{qQ}{r^2} \hat{r}_0$$



$$Q > q$$


$$F = F_2$$

$$-q$$

$$k = \frac{qQ}{(r-x)^2} = k \frac{qa}{x^2}$$

Contoh Soal

①.



$$q_3 = -2 \text{ C}$$

Diketahui : $q_1 = 1 \text{ C}$

$$q_2 = 3 \text{ C}$$

$$q_3 = -2 \text{ C}$$

Penyelesaian :

$$F_1 = F_2$$

$$k \frac{q_1 \cdot q_3}{x^2} = k \frac{q_3 \cdot q_2}{(10-x)^2}$$

$$k \frac{1 \cdot 2}{x^2} = k \frac{2 \cdot 3}{(10-x)^2}$$

$$\frac{1}{x^2} = \frac{3}{(10-x)^2}$$

$$(10-x)^2 = 3x^2$$

$$100 + x^2 - 20x = 3x^2$$

$$x^2 - 3x^2 - 20x = -100$$

$$-2x^2 - 20x = -100$$

$$-2x^2 - 20x + 100 = 0$$

$$-x^2 - 10x + 50 = 0$$

$$X_{(1,2)} = \frac{10 \pm \sqrt{b^2 - 4ac}}{2a}$$

$$= \frac{-10 \pm \sqrt{-10^2 - 4(-1)(50)}}{2 \cdot (-1)}$$

$$X_{(1,2)} = \frac{-10 \pm \sqrt{-10^2 - 4(-1)(50)}}{2 \cdot (-1)}$$

$$X_{(1,2)} = \frac{-10 \pm \sqrt{100 + 200}}{(-2)}$$


$$X_{(1,2)} = \frac{-10 \pm \sqrt{300}}{-2}$$

$$= \frac{-10 \pm \sqrt{\frac{300}{2}}}{2}$$

$$= \frac{-10 \pm 10\sqrt{3}}{2}$$

$$X_1 = \frac{10 + 17}{2} = \frac{27}{2} = 13,5$$

$$X_2 = \frac{10 - 17}{2} = \frac{-7}{2} = -3,5$$



$$q_3 = 1C$$

Diketahui : $q_1 = 1C$

$$q_2 = 2C$$

$$q_3 = 3C$$

Penyelesaian :

$$F_1 = F_2$$

$$k \frac{q_3 \cdot q_1}{(x^2)} = k \frac{q_3 \cdot q_2}{(10+x)^2}$$

$$\frac{1}{x^2} = \frac{2}{(10+x)^2}$$

$$-(10+x)^2 = -2x^2$$

$$2x^2 = 100 + x^2 + 20x$$

$$2x^2 - x^2 - 20x - 100 = 0$$

$$x^2 - 20x - 100 = 0$$

$$X_{(1,2)} = \frac{-b \pm \sqrt{-b^2 - 4ac}}{2 \cdot 1}$$


$$= \frac{-20 \pm \sqrt{(-20)^2 - (4 \cdot 1) 100}}{2}$$

$$= \frac{-20 \pm \sqrt{400 + 400}}{2}$$

$$= \frac{-20 \pm \sqrt{800}}{2}$$

$$= \frac{-20 \pm 20\sqrt{2}}{2} = \frac{-20 \pm 20\sqrt{2}}{2}$$

(3).



$$F_{12} = k \frac{q_1 \cdot q_2}{s^2} \left(\frac{-4\hat{i} + 3\hat{j}}{s} \right)$$

$$F_{13} = k \frac{q_1 q_3}{3^2} (-\hat{j})$$

$$\frac{(0\hat{i} - 3\hat{j})}{\sqrt{0^2 + 3^2}} = \frac{-3\hat{j}}{3} = -\hat{j}$$